

Florida Sharkman

Protocols

The information provided in this document does not, and is not intended to, constitute medical advice; instead, all information, content, and materials available in this document are for general informational purposes only. Information in this document may not constitute the most up-to-date medical or other information.

Readers of this document should contact their medical professionals to obtain advice with respect to any medical matter or condition. No reader of this document should act or refrain from acting on the basis of information in this document without first seeking medical advice from an accredited physician. Only your individual physician can provide assurances that the information contained herein — and your interpretation of it — is applicable or appropriate to your particular situation.

The views expressed in, or through, this document are those of the individual authors writing in their individual capacities only. All liability with respect to actions taken or not taken based on the contents of this document is hereby expressly disclaimed. The content on this posting is provided “as is;” no representations are made that the content is error-free.

Contents

Antiviral Protocol 3
Suggested Uses 3
Description 3
Timeframe 3
Antiviral Two-Month Calendar 4
Month One 4
Antiviral Protocol – Month Two 5
Protocol A 6
Protocol A – Monthly Calendar 6
Protocol B 7
Suggested Uses 7
Description 7
Timeframe 7
Protocol B – Monthly Calendar 8
Protocol C 9
Brain Cancer Glioblastoma (GBM) Protocol 10
Month One 10
Brain Cancer (GBM) Protocol – Month Two 11
Lyme Disease Protocol *(two-months)* 12
Month One 12
Lyme Disease Protocol – Month Two 13
Resources 14

Antiviral Protocol

Suggested Uses

- Viral infections such as: Covid, RSV, Epstein-Barr, HPV, etc.
- Viruses that cause many cancers:
    - HPV
    - Epstein-Barr
    - HCV
    - HIV
    - Cervical
    - HBV
    - HHV-8
    - HTLV-1
    - MCV
    - SV40
- Many gram-negative bacterial infections such as H. Pylori

Description

This protocol is based on the use of **Ivermectin**, **Doxycycline, Moxidectin**, **Fenbendazole**, and **Berberine**.

Timeframe

This is a two month-long protocol. If infection is cleared after this protocol is completed, no follow up is needed. If infection is not cleared, or for fighting virus-caused cancers, we recommend a two-week break and to proceed to Protocol B listed after Antiviral Protocol.

Antiviral Two-Month Calendar

Month One

Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Doxycycline 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg
Day 8	**Day 9**	**Day 10**	**Day 11**	**Day 12**	**Day 13**	**Day 14**
Doxycycline 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg
Day 15	**Day 16**	**Day 17**	**Day 18**	**Day 19**	**Day 20**	**Day 21**
Doxycycline 100 mg (AM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Fenbendazole** 50 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Fenbendazole** 50 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Fenbendazole** 50 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Fenbendazole** 50 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Fenbendazole** 50 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg
Day 22	**Day 23**	**Day 24**	**Day 25**	**Day 26**	**Day 27**	**Day 28**
Doxycycline 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg
Day 29	**Day 30**					
Doxycycline 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	Start Month Two. **NOTES:** **Berberine** – should be taken with food at lunch or dinner. **Doxycycline** – should be taken on an empty stomach with a large glass of water. If Doxycycline causes stomach upset, take it in conjunction with a light snack. **Fenbendazole** – take with the largest meal of the day. **Ivermectin** – take on an empty stomach.				

Antiviral Protocol – Month Two

Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Doxycycline 100 mg (AM) **Moxidectin** 0.5 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Berberine** 1200-1500 mg
Day 8	**Day 9**	**Day 10**	**Day 11**	**Day 12**	**Day 13**	**Day 14**
Doxycycline 100 mg (AM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Berberine** 1200-1500 mg
Day 15	**Day 16**	**Day 17**	**Day 18**	**Day 19**	**Day 20**	**Day 21**
Discontinue **Doxycycline** **Berberine** 1200-1500 mg	**Fenbendazole** 30 mg/kg **Berberine** 1200-1500 mg	**Fenbendazole** 30 mg/kg **Berberine** 1200-1500 mg	**Fenbendazole** 30 mg/kg **Berberine** 1200-1500 mg	**Fenbendazole** 30 mg/kg **Berberine** 1200-1500 mg	**Fenbendazole** 30 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg
Day 22	**Day 23**	**Day 24**	**Day 25**	**Day 26**	**Day 27**	**Day 28**
Ivermectin 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg
Day 29	**Day 30**					
Ivermectin 0.2 mg/kg **Berberine** 1200-1500 mg	**Moxidectin** 0.5 mg/kg **Berberine** 1200-1500 mg **COMPLETE**	If the infection is clear, congratulations, nothing further is required. If treating cancer or long-term infection, take a two-week break and proceed to Protocol B. **NOTES:** **Berberine** – should be taken with food at lunch or dinner. **Doxycycline** – should be taken on an empty stomach with a large glass of water. If Doxycycline causes stomach upset, take it in conjunction with a light snack. **Fenbendazole** – take a lower dose this month (30 mg/kg) with the largest meal. **Ivermectin** – take on an empty stomach.				

Protocol A

This is a general monthly protocol.

Protocol A – Monthly Calendar

Day 1	Day 2	*Full/New Moon* – Day 3	Day 4	Day 5	Day 6	Day 7
Doxycycline 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg
Day 8	**Day 9**	**Day 10**	**Day 11**	**Day 12**	**Day 13**	**Day 14**
Doxycycline 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg
Day 15	**Day 16**	**Day 17**	**Day 18** *New/Full Moon*	**Day 19**	**Day 20**	**Day 21**
Berberine 1200-1500 mg	**Berberine** 1200-1500 mg	**Oxfendazole** 20 mg/kg **Berberine** 1200-1500 mg	**Oxfendazole** 20 mg/kg **Berberine** 1200-1500 mg	**Oxfendazole** 20 mg/kg **Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg
Day 22	**Day 23**	**Day 24**	**Day 25**	**Day 26**	**Day 27**	**Day 28**
Berberine 1200-1500 mg	**Moxidectin** 0.5 mg/kg **Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg
Day 29	**Day 30**					
Berberine 1200-1500 mg	**Berberine** 1200-1500 mg	Repeat as many months as required to clear the condition. If Herxheimer Reactions are severe, reduce dosage or discontinue until the reactions have stopped. **NOTES:** **Berberine** – should be taken with food at lunch or dinner. **Doxycycline** – should be taken on an empty stomach with a large glass of water. If Doxycycline causes stomach upset, take it in conjunction with a light snack. **Fenbendazole** – take with the largest meal of the day. **Ivermectin** – take on an empty stomach. **Oxfendazole** – take with a fatty meal for better absorption.				

Protocol B

Suggested Uses

- Start Point for cancers other than viral
- Second phase for viral cancers
- Crohn's disease
- Colitis
- Ulcerated Colitis
- Parasitic worm infections
- Fungal infections
- Biofilms

Description

This protocol is based on the use of **Ivermectin**, **Doxycycline**, **Fenbendazole**, and **Berberine**.

Timeframe

This is a month-long protocol. If the infection is cleared after this protocol is completed, no follow up is needed. If the infection is not cleared, or for fighting cancers, this protocol can be used for many months until the condition clears.

Protocol B – Monthly Calendar

Day 1	Day 2	Day 3 *New/Full Moon*	Day 4	Day 5	Day 6	Day 7
Doxycycline 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg
Day 8	**Day 9**	**Day 10**	**Day 11**	**Day 12**	**Day 13**	**Day 14**
Doxycycline 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg
Day 15	**Day 16**	**Day 17**	**Day 18** *New/Full Moon*	**Day 19**	**Day 20**	**Day 21**
Berberine 1200-1500 mg	**Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg
Day 22	**Day 23**	**Day 24**	**Day 25**	**Day 26**	**Day 27**	**Day 28**
Ivermectin 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg
Day 29	**Day 30**					
Berberine 1200-1500 mg	**Berberine** 1200-1500 mg	Repeat as many months as required to clear the condition. **NOTES:** **Berberine** – should be taken with food at lunch or dinner. **Doxycycline** – should be taken on an empty stomach with a large glass of water. If Doxycycline causes stomach upset, take it in conjunction with a light snack. **Fenbendazole** – take with the largest meal of the day. **Ivermectin** – take on an empty stomach.				

Protocol C

This is a monthly protocol designed for maintenance, no sooner than after 6 months when parasites (nematodes) are no longer detected in the stool.

Day 1	Day 2	Day 3 *Full/New Moon*	Day 4	Day 5	Day 6	Day 7
Oxfendazole 20 mg/kg **Berberine** 1200-1500 mg	**Oxfendazole** 20 mg/kg **Berberine** 1200-1500 mg	**Oxfendazole** 20 mg/kg **Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg
Day 8	**Day 9**	**Day 10**	**Day 11**	**Day 12**	**Day 13**	**Day 14**
Berberine 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg
Day 15	**Day 16**	**Day 17**	**Day 18** *New/Full Moon*	**Day 19**	**Day 20**	**Day 21**
Berberine 1200-1500 mg	**Berberine** 1200-1500 mg	**Moxidectin** 0.5 mg/kg **Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg
Day 22	**Day 23**	**Day 24**	**Day 25**	**Day 26**	**Day 27**	**Day 28**
Berberine 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg	**Berberine** 1200-1500 mg
Day 29	**Day 30**					
Berberine 1200-1500 mg	**Berberine** 1200-1500 mg	Repeat as many months as required to clear the condition. **NOTES:** **Berberine** – should be taken with food at lunch or dinner. **Oxfendazole** – take with a fatty meal for better absorption.				

Brain Cancer Glioblastoma (GBM) Protocol

This is a 2-month (60 days) protocol.

Month One

Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Doxycycline 100 mg (AM) **Moxidectin** 0.5 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg
Day 8	**Day 9**	**Day 10**	**Day 11**	**Day 12**	**Day 13**	**Day 14**
Doxycycline 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg
Day 15	**Day 16**	**Day 17**	**Day 18**	**Day 19**	**Day 20**	**Day 21**
Fenbendazole 50 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Fenbendazole** 50 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Fenbendazole** 50 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Fenbendazole** 50 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Fenbendazole** 50 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg
Day 22	**Day 23**	**Day 24**	**Day 25**	**Day 26**	**Day 27**	**Day 28**
Ivermectin 0.2 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg
Day 29	**Day 30**					
Doxycycline 100 mg (AM) **Moxidectin** 0.5 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**NOTES:** **Berberine** – should be taken with food at lunch or dinner. **Doxycycline** – should be taken on an empty stomach with a large glass of water. If Doxycycline causes stomach upset, take it in conjunction with a light snack. **Fenbendazole** – take with the largest meal of the day. **Ivermectin** – take on an empty stomach.				

Brain Cancer (GBM) Protocol – Month Two

Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Doxycycline 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg
Day 8	**Day 9**	**Day 10**	**Day 11**	**Day 12**	**Day 13**	**Day 14**
Doxycycline 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Fenbendazole** 50 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Fenbendazole** 50 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg
Day 15	**Day 16**	**Day 17**	**Day 18**	**Day 19**	**Day 20**	**Day 21**
Fenbendazole 50 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Fenbendazole** 50 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Fenbendazole** 50 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg
Day 22	**Day 23**	**Day 24**	**Day 25**	**Day 26**	**Day 27**	**Day 28**
Ivermectin 0.2 mg/kg **Azithromycin** 500 mg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg
Day 29	**Day 30**					
Ivermectin 0.2 mg/kg **Berberine** 1200-1500 mg	**Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	Repeat as needed. **NOTES:** **Berberine** – should be taken with food at lunch or dinner. **Doxycycline** – should be taken on an empty stomach with a large glass of water. If Doxycycline causes stomach upset, take it in conjunction with a light snack. **Fenbendazole** – take with the largest meal of the day. **Ivermectin** – take on an empty stomach.				

Lyme Disease Protocol *(two-months)*

Month One

Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Doxycycline 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg
Day 8	**Day 9**	**Day 10**	**Day 11**	**Day 12**	**Day 13**	**Day 14**
Doxycycline 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg
Day 15	**Day 16**	**Day 17**	**Day 18**	**Day 19**	**Day 20**	**Day 21**
Doxycycline 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Doxycycline** 100 mg (PM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg
Day 22	**Day 23**	**Day 24**	**Day 25**	**Day 26**	**Day 27**	**Day 28**
Doxycycline 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg
Day 29	**Day 30**					
Doxycycline 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Doxycycline** 100 mg (PM) **Berberine** 1200-1500 mg					

NOTES:
Repeat if necessary with or without DOXY during Month Two.

Berberine – should be taken with food at lunch or dinner.
Doxycycline – should be taken on an empty stomach with a large glass of water.
If Doxycycline causes stomach upset, take it in conjunction with a light snack.
Fenbendazole – take with the largest meal of the day.
Ivermectin – take on an empty stomach.

Lyme Disease Protocol – Month Two

Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Doxycycline 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg
Day 8	**Day 9**	**Day 10**	**Day 11**	**Day 12**	**Day 13**	**Day 14**
Doxycycline 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg
Day 15	**Day 16**	**Day 17**	**Day 18**	**Day 19**	**Day 20**	**Day 21**
Doxycycline 100 mg (AM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Fenbendazole** 50 mg/kg (PM) **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg
Day 22	**Day 23**	**Day 24**	**Day 25**	**Day 26**	**Day 27**	**Day 28**
Doxycycline 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg
Day 29	**Day 30**					
Doxycycline 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**Doxycycline** 100 mg (AM) **Ivermectin** 0.2 mg/kg **Berberine** 1200-1500 mg	**NOTES:** **Low-dose DOXY during Month Two. If you are still having issues after Month Two, email info@floridasharkman.org** **Berberine** – should be taken with food at lunch or dinner. **Doxycycline** – should be taken on an empty stomach with a large glass of water. If Doxycycline causes stomach upset, take it in conjunction with a light snack. **Fenbendazole** – take with the largest meal of the day. **Ivermectin** – take on an empty stomach.				

Resources

Website: http://floridasharkman.org/

Truth Social: @FloridaSharkman